

**VIRGINIA RECREATIONAL FISHING DEVELOPMENT FUND
SUMMARY PROJECT APPLICATION***

NAME AND ADDRESS OF APPLICANT: Virginia Institute of Marine Science P. O. Box 1346 Gloucester Point, VA 23062-1346	PROJECT LEADER(name, phone, e-mail) Marcel Montane 804-684-7328 marcel@vims.edu
PRIORITY AREA OF CONCERN Tributaries of the Chesapeake Bay	PROJECT LOCATION Brackens and Wormley Ponds (York R.), Kamp's Millpond (Rappahannock R.) and Wareham's Pond (James R.)
DESCRIPTIVE TITLE OF PROJECT Estimating relative abundance of Young-of-Year American Eel, <i>Anguilla rostrata</i> , in the Virginia tributaries of Chesapeake Bay.	
PROJECT SUMMARY: The need for fisheries independent data from monitoring surveys is essential to many of the fishery management plans (FMP) for the Atlantic States Marine Fisheries Commission (ASMFC) and other management agencies. Specifically, this project meets the mandates of the ASMFC's FMP for American eel. Monitoring of glass eels (young-of-year) as they enter the estuary will provide estimates of recruitment in Virginia and allow for long-range planning for future harvestable stocks.	
EXPECTED BENEFITS: Recreational and commercial fishermen will benefit from this study as it will provide the Virginia Marine Resources Commission (VMRC) and ASMFC with an index of annual recruitment for juvenile American eels. The American eel is an important bait fishery in Virginia for game fish such as striped bass and cobia. Additionally, the American eel commercial fishery in Virginia from 2000-2006 landed an average of 116,458 lbs. Estimates of year class strength provide an "early warning" of recruitment success or failure, and are vital for proper species management.	
COSTS: VMRC Funding: \$45,348 Recipient Funding: \$7,256 TOTAL COST: \$52,604 Detailed budget included with proposal.	

Updated 6/1/05

*This form alone does not constitute a complete application, see application instructions or contact Sonya Davis at 757-247-8155 or sonya.davis@mrc.virginia.gov : Due dates are June 15 (Jul. - Nov. Cycle) and December 15 (Jan. - May Cycle)

Estimated Cost:

VMRC American Eel Budget 2008:

Estimating relative abundance of young of year American eel, *Anguilla rostrata*, in the Virginia tributaries of Chesapeake Bay.

Personnel	Time	MRFAB	VIMS	Total
Montane, PI	11%	6,274		6,274
Lowery	15%	5,488		5,488
Brooks	15%	5,475		5,475
Halvorson	15%	5,012		5,012
Rhea	5%	1,200		1,200
Fringe, 40% salaries		9,380		9,380
Supplies¹				
Field and lab supplies		1,000		1,000
Travel				
Field Sites		2,050		2,050
Regional Meeting (1 person)		400		400
Facilities & Administrative Costs		9,070	7,256	16,325
Totals		\$45,348	7,256	\$52,604

Facilities and Administrative Costs: F&A costs are assessed at 25% for funds provided by Marine Recreational Fishing Advisory Board. Due to the critical nature of the funding shortfall, VIMS will provide a majority of the F&A costs associated with this project.

¹Supplies include waders, water temperature dataloggers, materials for eel traps, waterproof paper, etc.

**Estimating relative abundance of Young-of-Year American Eel,
Anguilla rostrata, in the Virginia tributaries of the Chesapeake Bay.**

Proposal Submitted to:

Virginia Marine Resources Commission
Marine Recreational Fishing Advisory Board
2400 Washington Avenue, Newport News, VA 23607


15 January 2008 – 31 December 2008

Submitted by:

Virginia Institute of Marine Science, School of Marine Science
College of William and Mary, Gloucester Point, VA, 23062




Marcel Montane, Principal Investigator
Marine Scientist Supervisor
Dept. of Fisheries Science



John Olney
Chair, Dept. of Fisheries Science



Roger Mann
Director for Research and Advisory Services



Jane Lopez
Director of Sponsored Research

December 5, 2007

Title: Estimating relative abundance of the young-of-year American eel, *Anguilla rostrata*, in the Virginia tributaries of Chesapeake Bay.

Introduction

The Atlantic States Marine Fisheries Commission (ASMFC) adopted an interstate fishery management plan for American eel (*Anguilla rostrata*) in 1999 (ASMFC, 2000). The plan mandated that all states/jurisdictions would be required to conduct an annual fishery-independent survey for young-of-year (YOY) American eel using specific methods and gears approved by the ASMFC Technical Committee. These surveys would be required of all states/jurisdictions beginning in 2001. The importance of the American eel commercial fishery in Virginia is well known. Additionally, this fishery also produces a large bait component for recreational anglers with little known about the potential impact bait harvest has on the overall fishery, or the eel population in general.

In Virginia, during Spring 2000, exploratory surveys were conducted by the Virginia Institute of Marine Science (VIMS) to establish appropriate sampling gear and methodologies to evaluate YOY American eel recruitment. Since 2001, both the VMRC Marine Recreational Fishing Advisory Board and the Commercial Fishing Advisory Board have supported this project.

History and Relevance

VIMS has a long history of monitoring and assessing the fishery stocks in the lower Chesapeake Bay. In particular, there are several long-term programs specifically targeting juvenile fishes with a primary goal of obtaining annual recruitment estimates (i.e. VIMS Trawl Survey since 1955 and the VIMS Striped Bass Seine Survey, initiated in 1967). Although American eel are often captured by these surveys, most are juveniles (yellow eels, total length or TL > 180 mm), with very few YOY (glass eels) collected. Only a few states/jurisdictions (Maine, Maryland, New Jersey, and Nova Scotia, Canada) have historically collected information on glass eel recruitment. With the new ASMFC mandate, a coastwide estimate of annual recruitment for glass eels was established.

Information for American eel is often limited or lacking possibly due to its unique life history. The American eel is a catadromous species, present along the Atlantic and Gulf coasts of North America and inland in the St. Lawrence Seaway and Great Lakes (Murphy et al., 1997). The species is panmictic and supported throughout its range by a single spawning population (Haro et al., 2000). Spawning takes place during winter to early spring in the Sargasso Sea. The eggs hatch into leaf-shaped ribbon-like larvae called leptocephali, which are transported by the ocean currents (over 9-12 months) in a generally northwesterly direction. Within a year, metamorphosis into the next life stage (glass eel) occurs in the Western Atlantic near the East Coast of North America. Coastal currents and active migration transport the glass eels into rivers and estuaries from February to June in Virginia and Maryland. As growth continues, the eel becomes pigmented (elver stage) and within 12 –14 months acquires a dark color with underlying yellow (yellow eel stage; Facey and Van Den Avyle, 1987). Many eels migrate upriver into

freshwater rivers, streams, lakes, and ponds, while others remain in estuaries. Most of the eel's life is spent in these habitats as a yellow eel. Age at maturity varies greatly with location and latitude, and in Chesapeake Bay may range from 8 to 24 years, with most eels less than ten years old (Owens and Geer, 2003). *A. rostrata* from Chesapeake Bay mature and migrate at an earlier age than eels from northern areas (Hedgepeth, 1983). Upon maturity, eels migrate back to the Sargasso Sea to spawn and die (Haro et al., 2000). Metamorphosis into the silver eel stage occurs during the seaward migration that occurs from late summer through autumn.

It has been suggested that glass eel migration consists of waves of invasion (Boetius and Boetius, 1989 as reported by Ciccotti et al., 1995), and perhaps a fortnightly periodicity related to selective tidal stream transport (Ciccotti et al., 1995). Additionally alterations in freshwater inflow (patterns and magnitudes) to bays and estuaries may alter flow regimes and consequently affect the size, timing and spatial patterns of upstream migration of glass eels and elvers (Facey and Van Den Avyle, 1987).

The American eel is a valuable commercial species along the entire Atlantic Coast from New Brunswick to Florida, with Virginia historically accounting for one-third of annual Atlantic and Gulf Coast landings and Chesapeake Bay one-half of the total (NMFS, 2006). Landings along the U.S. Atlantic Coast have varied from 290 MT in 1962 to a high of 1600 MT in 1975 (NMFS, 2006) and the American eel once accounted for more than 25% of the total fish biomass of East Coast streams (ASMFC, 2005).

In recent years there has been a decline in landings in Virginia with similar patterns seen in the Canadian Maritime Provinces. These declines are also apparent in fishery-independent trawl surveys in Chesapeake Bay (Fabrizio and Montane, 2007). Although primarily a commercial species, an unknown portion of the total harvest in Virginia is part of the bait fishery for coastal gamefish (Jenkins and Burkhead, 1993) such as striped bass and cobia. Information available from the National Marine Fisheries Service (NMFS) regarding Atlantic Coast eel harvest for bait also indicates a large decline in harvest since the mid-1980's. As these recreational fisheries continue to grow, the amount of eels sold as bait to support the fishery will grow as well.

American eel are not usually considered a sport fish, though they may be caught by recreational fishermen (Collette and Klein-MacPhee, 2002). The NMFS Marine Recreational Fisheries Statistics Survey shows a similar declining trend in catch of eels for Virginia through 2006 (USDOC, 2006).

With coastwide declines in harvest observed throughout the Atlantic States and Maritime Provinces, it is essential that reliable information on recruitment and other life history parameters (e.g. length and age, age at maturity) be obtained to establish a reliable stock assessment plan. Hypotheses for the coastwide decline include shifts in the Gulf Stream which affect recruitment, pollution, overfishing, parasites which affect migration and spawning, and up-river impediments, such as fish barriers (Castonguay et al., 1994).

Many present fisheries management techniques cannot be applied to American eels because basic data on its biology are either missing or studies have shown conflicting results. Different growth rates between water systems, and large variations for length at age have made it difficult to perform stock assessments (Owens and Geer, 2003). Additionally, few studies have addressed the recruitment of glass eels to Atlantic Coast estuaries from the spawning grounds in

the Sargasso Sea, and long-term datasets for American eels are lacking (Powles and Warlen, 2002).

Need:

Measures of juvenile abundance are widely employed as a key element in the management of many Atlantic states coastal fishery resources. Fluctuations in relative abundance of early juveniles (age 0 or YOY) generated from fisheries-independent survey programs have been found to provide a reliable and early estimator of future year class strength for species such as striped bass (Goodyear, 1985) and crabs (Lipcius and Van Engel, 1990) in Chesapeake Bay. For example, the current Interstate Fisheries Management Plan for the striped bass relies heavily on estimates of juvenile abundance, both as 'action levels' for the intensification and relaxation of restrictions and as a measure of year class strength in mathematical population models (USDOI and USDOC, 1989). In addition to providing FMP input, juvenile indices can be an "early warning" of year class failure. ASMFC mandates that all Atlantic coastal states and jurisdictions conduct an annual YOY American eel survey.

Objectives:

1. To monitor the glass eel migration, or run, into the Virginia Chesapeake Bay tributaries, to determine spatial and temporal components of recruitment.
2. Examine the tidal, lunar, and hydrographic parameters (temperature, pH, etc.) which may influence glass eel recruitment.
3. Collect basic biological information on glass eels (i.e. length, weight, and pigment stage).

Field work will be performed from late February 2008 to late May 2008 likely corresponding to the period of maximum YOY American eel recruitment.

Expected Benefits and Results:

A primary benefit of this project is to insure that recruitment of YOY American eel will be monitored in the tidal waters of Virginia. Since the Chesapeake Bay jurisdictions (Virginia, Maryland, and the Potomac River Fisheries Commission) comprise nearly 63% of the commercial landings on the Atlantic East Coast, monitoring annual recruitment in this region constitutes a key element in multi-state efforts to manage this Atlantic coastal fishery resource. This need was further emphasized by the Atlantic Coastal Cooperative Fishery Management Act (PL-103-206), specifying that states identified in ASMFC management must be in compliance, with the American Eel FMP directly requiring monitoring of YOY recruitment.

The information collected from this study will provide updated information on the timing and distribution of YOY American eel recruitment into the Chesapeake Bay's tributaries. The

effects of tidal and lunar factors on abundance will be examined as well as relationships between abundance and hydrographic parameters. The information collected will be used by resource management agencies on the State, Federal, and possibly International levels, to better understand recruitment of this catadromous species and provide a valuable tool for assessing the success of present management strategies. Virginia YOY eel survey data was incorporated into the recent ASMFC Update of the American Eel Stock Assessment Report (ASMFC, 2006). In this report, the ASMFC Peer Review Panel reinforced the importance of the coastwide YOY eel survey as the surveys could serve as “an index of sustained recruitment over the historical coastal range, an early warning of recruitment failure, and/or an early warning of potential range contraction of the species (ASMFC, 2006)”.

Approach:

Minimum criteria for YOY American eel sampling has been proposed by the ASMFC American Eel Technical Committee, and approved by the American Eel Management Board. Due to the importance of the eel fishery in Virginia, additional sites are sampled to insure proper temporal and spatial coverage, and to provide reliable recruitment estimates.

To provide the necessary spatial coverage at least one site should be sampled per major tributary (the James, York, and Rappahannock Rivers). Site selection will be based on known areas of glass eel migrations, accessibility, and specific criteria which have been documented as sources of glass eel concentration (such as millponds which feed directly into the estuaries). Prior sampling has revealed several sites that met these criteria and produced adequate numbers of glass eels (Montane et al., 2007). Current (and proposed) sites to be sampled include Brackens and Wormley Ponds on the York River, Kamp’s Millpond on the Rappahannock River and Wareham’s Pond on the James River (Figure 1).

The minimum sampling period based on ASMFC’s mandates is four days per week for at least six weeks. Previous years results indicate the run in Virginia to be more protracted, requiring a longer sampling period (Montane et al., 2007). In accordance with the ASMFC mandate, sites will be sampled a minimum of four days per week (usually Monday, Tuesday, Thursday and Friday), starting around end of February/early March. Sampling will continue throughout the glass eel run (end of May/early June) or until 0 glass eels are caught. To determine the start of the run, sites will be sampled at least once weekly until glass eels recruit to the eel trap, indicating the start of the sampling period. When catches drop substantially later in the season, sampling effort will be reduced until the run is completed. Over several years, it is hoped that the timing of the run may become more predictable and this window of preliminary sampling might be diminished, though the start (and end) of the run has been variable between years.

At each site (sites may be combined if on the same tributary) a sample of sixty eels will be collected weekly, measured to the nearest 0.1mm TL, weighed to the nearest 0.01g, and pigment stage recorded (see Haro and Krueger, 1988 for staging criteria). The remaining catch will be enumerated and placed above the spillway. At each site temperature, tidal stage, time, and condition of the gear will be recorded. The gear utilized will be an Irish eel ramp (B. Jessops design; see Montane et al., 2007 for gear configuration).

Sites chosen in the past have proven to be excellent locations to study YOY eel recruitment and these sites will again be studied to lengthen the time series. Brief results from the past surveys follow. Sampling at the York River sites (i.e., Brackens Pond and Wormley Pond, see Figure 1) was conducted from 14 February to 18 May 2007. Wareham's Pond on the James River (see Figure 1) was sampled from 15 February to 18 May 2007, while sampling at Kamp's Millpond on the Rappahannock River (see Figure 1) was conducted from 27 February 2007 through 5 July 2007. Glass eel and elver CPUE's at each site were standardized to a 24 hour soak time for the Irish eel ramp, and geometric means were calculated using the time period in which 95% of the cumulative total catch was sampled (i.e. dates in which 0% - 2.5% and 97.5% - 100% of the cumulative total catch were excluded), in an effort to account for the interannual variability in the period of maximum recruitment.

The CPUEs for both glass eels and elvers have been variable at all sites over the time series. The 2007 glass eel CPUE decreased at both Brackens Pond and Wormley Pond compared to 2006 (Figure 2, top), while the elver CPUE for 2007 increased at both sites relative to the previous year (Figure 2, bottom). Kamp's Millpond 2007 glass eel CPUE increased very slightly compared to 2006, while CPUE decreased at Wareham's Pond in 2007 (Figure 3, top). Kamp's Millpond 2007 elver CPUE decreased relative to the 2006 value (Figure 3, bottom). Conversely, the elver CPUE increased at Wareham's Pond in 2007 compared to 2006 (Figure 3, bottom).

Location:

Proposed sites to be sampled include Brackens and Wormley Ponds on the York River, Kamp's Millpond on the Rappahannock River and Wareham's Pond on the James River (see Figure 1).

Deliverables and Final Products:

Quarterly progress reports and a final report will be submitted to VMRC MRFAB according to their reporting requirements. Data will be made available to ASMFC to fulfill their mandate and also presented at local, regional and national/international meetings as well as submitted for publication in scientific journals.

REFERENCES

- ASMFC, 2000. Interstate Fishery Management Plan for American eel, *Anguilla rostrata*. 79 pp.
- ASMFC, 2005. Addendum I to the Interstate Fishery Management Plan for American eel. 11pp.
- ASMFC, 2006. Update of the American Eel Stock Assessment Report. American Eel Stock Assessment Subcommittee and the American Eel technical Committee. 51 pp.
- Castonguay, M. P., P. V. Hodson, M. J. Eckersley, J-D. Dutil and G. Vereault. 1994. Why is recruitment of the American eel, *Anguilla rostrata*, declining in the St. Lawrence River and Gulf? *Can. J. Fish. Aquat. Sci* 51:479-488.
- Ciccotti, E., T. Ricci, M. Scardi, E. Fresi and S. Cataudella. 1995. Intra-seasonal characterization of glass eel migration in the River Tiber: space and time dynamics. *J. Fish Biol.* 47:248-255.
- Collette, B. B. and G. Klein-MacPhee. 2002. *Bigelow and Schroeders Fishes of the Gulf of Maine*. 3rd Ed. Smithsonian Institution Press. Washington, D.C. 748 pp.
- Fabrizio, M.C and M. M. Montane. 2007. Estimating relative juvenile abundance of ecologically important finfish and invertebrates in the Virginia portion of Chesapeake Bay. Report to the NOAA Chesapeake Bay Office, Award No. NA03NMF4570378. Virginia Institute of Marine Science, Gloucester Point, VA. 97 pp.
- Facey, D. E. and M. J. Van Den Avyle. 1987. Species profiles: life histories and environmental requirements of coastal fishes and invertebrates (North Atlantic)—American eel. U. S. Fish Wildl. Serv. Biol. Rep. 82(11.74). U. S. Army Corps of Engineers, TR EL-82-4. 28 pp.
- Goodyear, C. P. 1985. Relationship between reported commercial landings and abundance of young striped bass in Chesapeake Bay, Maryland. *Trans. Amer. Fish. Soc.* 114(1):92-96.
- Haro, A.J. and W.H. Krueger. 1988. Pigmentation, size, and migration of elvers (*Anguilla rostrata* (Lesueur)) in a coastal Rhode Island stream. *Can. J. Zool.* 66: 2528-2533.
- Haro, A., W. Richkus, K. Whalen, W.-Dieter Busch, S. Lary, T. Brush, and D. Dixon. 2000. Population decline of the American eel: Implications for Research and management. *Fisheries* 25(9): 7-16.
- Hedgepeth, M. Y. 1983. Age, growth and reproduction of American eels, *Anguilla rostrata* (Lesueur), from the Chesapeake Bay area. Masters Thesis. College of William and Mary. 61 pp.
- Jenkins, R. E. and N. M. Burkhead. 1993. *Freshwater fishes of Virginia*. American Fisheries Society. Bethesda, MD. 1079 pp.
- Lipcius, R. N. and W. A. Van Engel. 1990. Blue crab population dynamics in Chesapeake Bay: variation in abundance (York River, 1972-1988) and stock-recruit functions. *Bull. Mar. Sci.* 46(1): 180-194.
- Montane, M.M., W.A. Lowery, H. Brooks and A.D. Halvorson. 2007. Estimating Relative Abundance of Young of Year American Eel, *Anguilla rostrata*, in the Virginia Tributaries of Chesapeake Bay (Spring 2006). Final Report to the Virginia Marine Resources Commission. Project # RF/CF06-01. 32 pp.
- Murdy, E.O., R.S. Birdsong and J.A. Musick. 1997. *Fishes of Chesapeake Bay*. Smithsonian

Institution Press. 324 pp.

NMFS, 2006. "Annual commercial landings statistics. National Marine Fisheries Service Fisheries Statistics Division Annual Landings Query".
 <http://remora.ssp.nmfs.gov/MFPUBLIC/owa/mrfss.FT_HELP.SPECIES>.

Owens, S. J. and P. J. Geer. 2003. Size and age of American eels collected from tributaries of the Virginia portion of the Chesapeake Bay. *American Fisheries Society Symposium Series* 33:117-124.

Powles, P. M. and S. M. Warlen. 2002. Recruitment season, size and age of young American eels (*Anguilla rostrata*) entering an estuary near Beaufort, North Carolina. *Fish. Bull.* 100:299-306.

USDOI and USDOC (United States Department of the Interior and United States Department of Commerce). 1989. Emergency striped bass research study. Report for 1988. Washington, DC, USA. 55p.

USDOC (United States Department of Commerce), 2006. "Marine recreational fisheries statistics survey catch snapshot query".
http://remora.ssp.nmfs.gov/mrfss/queries/catch_snapshot.html

Figure 1. 2007 YOY Eel Sampling Sites.




Figure 2. YOY (top) and Elver (bottom) CPUE (Geometric Means) for Brackens (2000-2007) and Wormley Ponds (2001-2007).





Figure 3. YOY (top) and Elver (bottom) CPUE (Geometric Means) for Kamps (2000-2007) and Warehams Ponds combined (2003-2007).

